

Functional Callan-Symanzik equation for QED

J. Alexandre¹, J. Polonyi^{2,3}, and K. Sailer⁴

¹ Department of Physics, King's College, London, UK

² Institute for Theoretical Physics, Louis Pasteur University, Strasbourg,
France

³ Department of Atomic Physics, Lorand Eötvös University Budapest,
Hungary

⁴ Department for Theoretical Physics, University of Debrecen, Debrecen,
Hungary

Abstract

An exact evolution equation, the functional generalization of the Callan-Symanzik method, is given for the effective action of QED where the electron mass is used to turn the quantum fluctuations on gradually. The usual renormalization group equations are recovered in the leading order but no Landau pole appears.

Introduction: The idea of dimensional transmutation [1] is that a dimensionless parameter is traded for a dimensionful one. This replacement can be used to generate a renormalization flow where the field amplitude which is a quantity related to the size of the quantum fluctuations is evolved. These flows ("fluctuation flows") prove to be equivalent with the usual momentum flows at one-loop order. This scheme is developed in this paper for QED by constructing the evolution of the one-particle irreducible (1PI) generator functional where a mass parameter controls the quantum fluctuations. We derive an exact equation describing this evolution and will recover the usual one-loop momentum flows. We conclude with some observations on the connection between the two renormalization schemes.

The functional approach to renormalization group started with the use of a 'sharp cut-off' k , i.e. the elimination of the Fourier modes with $|p| > k$ was performed [2]. An infinitesimal change $k \rightarrow k + \Delta k$ produces the renormalization group (RG) equation for the Wilsonian, blocked action. This procedure has been developed later [3], as well as other schemes involving a 'smooth cut-off', i.e. where the Fourier modes are suppressed by means of a smooth, regulated version of the step-function and the evolution of the effective action is sought [4, 5]. The work presented here, although being quite different from a formal point of view, is actually motivated by the smooth cut-off scheme for the evolution of the effective action. The difference compared to that method is that the suppression depends on the amplitude rather than the characteristic wave-length of the modes. This method has been developed for a scalar model [6] where the usual RG equations were recovered in the U.V. regime, i.e. where the mass is close to the cut-off. Finally, the same functional method was also used in QED with an external field [7] where the evolution of the generator functional of the one-particle irreducible (1PI) graphs with the amplitude of the external field led to the dependence of the 1PI graphs on the external gauge.

Evolution equation: We start with the following bare Lagrangian in dimension $d = 4 - \varepsilon$

$$\mathcal{L} = \frac{1}{2e^2\mu^\varepsilon} \mathcal{A}_\mu \square (T^{\mu\nu} + \alpha L^{\mu\nu}) \mathcal{A}_\nu + \overline{\Psi} (i \not{\partial} - \not{\mathcal{A}} - zm_0) \Psi, \quad (1)$$

where $T^{\mu\nu}$ and $L^{\mu\nu}$ are respectively the transverse and longitudinal projectors in the inverse photon propagator, and the gauge parameter α characterizes the gauge fixing. The parameter z is introduced to control the amplitude of the fluctuations. For $z \gg 1$ the theory is dominated by a free mass term contribution and is perturbative. As z decreases the interaction with the gauge field becomes more important and quantum corrections increase in amplitude. Our aim is to study the evolution in z of the generator functional Γ_z of the 1PI diagrams, the effective action. The functional W_z of the connected diagrams is given by

$$\begin{aligned} & \exp W_z[\overline{\eta}, \eta, j^\mu] \\ &= \int \mathcal{D}[\overline{\Psi}, \Psi, \mathcal{A}_\mu] \exp \left\{ i \int_x \mathcal{L} + i \int_x (j^\mu \mathcal{A}_\mu + \overline{\eta} \Psi + \overline{\Psi} \eta) \right\}, \end{aligned} \quad (2)$$

and has the following functional derivatives

$$\begin{aligned}
\frac{\delta}{\delta \bar{\eta}} W &= i \langle \Psi \rangle = i\psi, \\
W \frac{\delta}{\delta \eta} &= i \langle \bar{\Psi} \rangle = i\bar{\psi}, \\
\frac{\delta}{\delta j_\mu} W &= i \langle \mathcal{A}^\mu \rangle = iA^\mu.
\end{aligned} \tag{3}$$

The effective action $\Gamma_z[\bar{\psi}, \psi, A_\mu]$ is defined as the Legendre transform of W_z ,

$$W_z = i\Gamma_z + i \int_x \left(j^\mu A_\mu + \bar{\eta}\psi + \bar{\psi}\eta \right), \tag{4}$$

and has the following functional derivatives

$$\frac{\delta}{\delta \bar{\psi}} \Gamma = -\eta, \quad \Gamma \frac{\delta}{\delta \psi} = -\bar{\eta}, \quad \frac{\delta}{\delta A_\mu} \Gamma = -j^\mu. \tag{5}$$

The functional manipulations which lead to the evolution of Γ_z are the same as those discussed in [7]: we first remark that the derivatives (3) and (5) of the functionals W_z and Γ_z with respect to their variables imply that

$$\begin{aligned}
\partial_z \Gamma &= -i\partial_z W = \left\langle -m_0 \int_x \bar{\Psi} \Psi \right\rangle \\
&= -m_0 \int_x \bar{\psi} \psi - m_0 \int_x \frac{\delta}{\delta \bar{\eta}} W \frac{\delta}{\delta \eta}.
\end{aligned} \tag{6}$$

Then the relation

$$\frac{\delta}{\delta \bar{\psi}} \Gamma \frac{\delta}{\delta \psi} = -i \left(\frac{\delta}{\delta \bar{\eta}} W \frac{\delta}{\delta \eta} \right)^{-1} \tag{7}$$

leads us to the final equation for the generator functional of the 1PI graphs,

$$\partial_z \Gamma_z + m_0 \int_x \bar{\psi} \psi = im_0 \text{Tr} \left\{ \left(\frac{\delta \Gamma}{\delta \bar{\psi} \delta \psi} \right)^{-1} \right\}, \tag{8}$$

where 'Tr' denotes the trace over space-time and Dirac indices. The inverse matrix $\delta^2 \Gamma^{-1}$ has to be taken with respect to the field variables, space-time and Dirac indices. We stress that Eq.(8) is exact and does not make reference

to any truncation of any sort. Note the similarity between Eq.(8) and other RG equations [5]. This similarity reflects the qualitative agreement between the usual momentum flows and our method, as we will discuss at the end of this paper.

Gradient expansion: In order to find an approximate solution of the evolution equation, we make an expansion in the amplitude and the gradient of the fields and use the ansatz

$$\Gamma_z[\bar{\psi}, \psi, A_\mu] = \int_x \left[A_\mu(x) \mathcal{D}^{-1\mu\nu} A_\nu(x) + \bar{\psi}(x) \mathcal{G}^{-1} \psi(x) - Z(z) \bar{\psi}(x) \not{A}(x) \psi(x) \right], \quad (9)$$

where the photon and fermion propagators are

$$\begin{aligned} \mathcal{D}^{-1\mu\nu}(p) &= \frac{\beta_T(z)}{e^2 \mu^\varepsilon} p^2 T^{\mu\nu}(p) + \alpha \frac{\beta_L(z)}{e^2 \mu^\varepsilon} p^2 L^{\mu\nu}(p), \\ \mathcal{G}^{-1}(p) &= Z(z) \not{p} - m(z) - z m_0, \end{aligned} \quad (10)$$

with $T^{\mu\nu}(p) = g^{\mu\nu} - p^\mu p^\nu / p^2$ and $L^{\mu\nu}(p) = p^\mu p^\nu / p^2$. The evolution of $\beta_T(z)$, $\beta_L(z)$, $Z(z)$ and $m(z)$ are obtained by expanding both sides of Eq.(8) in powers of the gauge field and by identifying the operators on the left and right hand sides. For this end we need the inverse of the matrix $\Gamma^{(2)}$

$$\Gamma^{(2)} = \begin{pmatrix} \frac{\delta\Gamma}{\delta\bar{\psi}\delta\psi} & \frac{\delta^2\Gamma}{\delta\bar{\psi}\delta\bar{\psi}} & \frac{\delta^2\Gamma}{\delta A_\mu\delta\bar{\psi}} \\ \frac{\delta\Gamma}{\delta\psi\delta\bar{\psi}} & \frac{\delta\Gamma}{\delta\psi\delta\psi} & \frac{\delta\Gamma}{\delta A_\mu\delta\psi} \\ \frac{\delta\Gamma}{\delta\psi\delta A_\mu} & \frac{\delta^2\Gamma}{\delta\psi\delta A_\mu} & \frac{\delta^2\Gamma}{\delta A_\mu\delta A_\nu} \end{pmatrix}, \quad (11)$$

which will be obtained by expanding in the diagonal part $\Gamma_\Delta^{(2)}$ in momentum space,

$$\begin{aligned} (\Gamma^{(2)})^{-1} &= (\Gamma_\Delta^{(2)})^{-1} - (\Gamma_\Delta^{(2)})^{-1} \Gamma_{nd}^{(2)} (\Gamma_\Delta^{(2)})^{-1} \\ &+ (\Gamma_\Delta^{(2)})^{-1} \Gamma_{nd}^{(2)} (\Gamma_\Delta^{(2)})^{-1} \Gamma_{nd}^{(2)} (\Gamma_\Delta^{(2)})^{-1} \\ &- (\Gamma_\Delta^{(2)})^{-1} \Gamma_{nd}^{(2)} (\Gamma_\Delta^{(2)})^{-1} \Gamma_{nd}^{(2)} (\Gamma_\Delta^{(2)})^{-1} \Gamma_{nd}^{(2)} (\Gamma_\Delta^{(2)})^{-1} + \dots, \end{aligned} \quad (12)$$

where $\Gamma_{nd}^{(2)} = \Gamma^{(2)} - \Gamma_\Delta^{(2)}$. We obtain in this manner the result

$$\begin{aligned} \left(\frac{\delta\Gamma}{\delta\bar{\psi}\delta\psi} \right)^{-1} &= \mathcal{G} + \mathcal{G} \left(-c + c\mathcal{G}c - c\mathcal{G}c\mathcal{G}c + a^\mu \mathcal{D}_{\mu\nu} \tilde{b}^\nu \right. \\ &\quad \left. - a^\mu \mathcal{D}_{\mu\nu} \tilde{b}^\nu \mathcal{G}c - c\mathcal{G}a^\mu \mathcal{D}_{\mu\nu} \tilde{b}^\nu \right) \mathcal{G} + \dots, \end{aligned} \quad (13)$$

where

$$\begin{aligned}
a^\mu(p, q) &= -Z(z)\gamma^\mu\psi(-p-q), \\
b^\nu(p, q) &= \bar{\psi}(-p-q)Z(z)\gamma^\nu, \\
c(p, q) &= -Z(z)\tilde{A}(-p-q),
\end{aligned} \tag{14}$$

and the tilde stands for the transposed in momentum and spinor spaces. The evolution equations read finally as

$$\begin{aligned}
\partial_z \mathcal{D}^{-1\mu\nu}(p) &= -2im_0 Z^2(z) \int_q \text{tr} \left\{ \mathcal{G}^2(q) \gamma^\mu \mathcal{G}(p+q) \gamma^\nu \right\}, \\
\partial_z \mathcal{G}^{-1}(p) + m_0 &= -im_0 Z^2(z) \int_q \mathcal{D}_{\mu\nu}(q-p) \gamma^\mu \mathcal{G}^2(q) \gamma^\nu, \\
\partial_z Z(z) \gamma^\mu &= -2m_0 Z^3(z) \int_q \mathcal{D}_{\kappa\nu}(q) \gamma^\nu \mathcal{G}^2(q) \gamma^\mu \mathcal{G}(q) \gamma^\kappa.
\end{aligned} \tag{15}$$

The following remarks are important:

- The photon does not acquire a mass in the evolution in z and the IR divergences are the usual ones.
- The longitudinal part of the photon propagator does not evolve with z . Accordingly, the fluctuations do not generate longitudinal contributions to photons.

Furthermore a technical point, it is the evolution equation of the fermion mass only which needs regulator, the other equations are divergence-free with the given effective action ansatz.

To compare this result with the traditional method based on the loop-expansion, note that the change $zm_0 \rightarrow (z + \delta z)m_0$ of the bare electron mass changes the fermion propagator as $\mathcal{G} \rightarrow \mathcal{G} + \mathcal{G}\delta zm_0\mathcal{G}$ in the internal lines of the Feynman graphs. This observation, the starting point of the Callan-Symanzik method is sufficient to realize that the evolution equations (15) can obviously be written as

$$\begin{aligned}
\partial_z \mathcal{D}^{-1\mu\nu}(p) &= -i\partial_z \int_q \text{tr} \left\{ \mathcal{G}(q) \gamma^\mu \mathcal{G}(p+q) \gamma^\nu \right\} + \mathcal{O}(\hbar^2), \\
\partial_z \mathcal{G}^{-1}(p) + m_0 &= -i\partial_z \int_q \mathcal{D}_{\mu\nu}(q-p) \gamma^\mu \mathcal{G}(q) \gamma^\nu + \mathcal{O}(\hbar^2), \\
\partial_z Z \gamma^\mu &= -\partial_z \int_q \mathcal{D}_{\kappa\nu}(q) \gamma^\nu \mathcal{G}(q) \gamma^\mu \mathcal{G}(q) \gamma^\kappa + \mathcal{O}(\hbar^2),
\end{aligned} \tag{16}$$

whose integrals give immediately

$$\begin{aligned}
\mathcal{D}^{-1\mu\nu}(p) &= \mathcal{D}_{tree}^{-1\mu\nu}(p) - i \int_q tr \{ \mathcal{G}(q) \gamma^\mu \mathcal{G}(p+q) \gamma^\nu \} + \mathcal{O}(\hbar^2), \\
\mathcal{G}^{-1}(p) &= \mathcal{G}_{tree}^{-1}(p) - i \int_q \mathcal{D}_{\mu\nu}(q-p) \gamma^\mu \mathcal{G}(q) \gamma^\nu + \mathcal{O}(\hbar^2), \\
Z\gamma^\mu &= Z_{tree}\gamma^\mu - \int_q \mathcal{D}_{\kappa\nu}(q) \gamma^\nu \mathcal{G}(q) \gamma^\mu \mathcal{G}(q) \gamma^\kappa + \mathcal{O}(\hbar^2),
\end{aligned} \tag{17}$$

the usual one-loop corrections to the 1PI functions appearing in the ansatz (9).

Beta functions: The computation of the integrals is straightforward and leads to the following result,

$$\begin{aligned}
\beta'_L(z) &= 0, \\
\beta'_T(z) &= \frac{e^2 \mu^\varepsilon}{6\pi^2} \frac{m_0}{zm_0 + m(z)}, \\
Z'(z) &= \frac{e^2 \mu^\varepsilon}{8\pi^2 \alpha} \frac{Z(z)m_0}{zm_0 + m(z)}, \\
m'(z) &= -\frac{m_0}{8\pi^2} \frac{e^2 \mu^\varepsilon}{\varepsilon} \left(\frac{3}{\beta_T(z)} + \frac{1}{\alpha} \right) + \text{finite},
\end{aligned} \tag{18}$$

where the prime denotes a derivative with respect to z . In what follows, we will keep the bare theory and $\varepsilon \neq 0$. We define $m(z) = m_0 \phi(z)$ and write $\beta(z) = \beta_T(z)$, such that the equations (18) read

$$\begin{aligned}
\phi'(z) &= -\frac{e^2 \mu^\varepsilon}{8\pi^2 \varepsilon} \left(\frac{3}{\beta(z)} + \frac{1}{\alpha} \right), \\
\beta'(z) &= \frac{e^2 \mu^\varepsilon}{6\pi^2} \frac{1}{z + \phi(z)}, \\
Z'(z) &= \frac{e^2 \mu^\varepsilon}{8\pi^2 \alpha} \frac{Z(z)}{z + \phi(z)}.
\end{aligned} \tag{19}$$

In order to compare these scaling laws with the one-loop result, we look at the situation of weak fluctuations where the fermion mass term is dominant in the bare action, i.e. where z is close to some initial value $z_0 \gg 1$ and the parameter $m(z)$ is close to zero. In this regime we have $z + \phi(z) \simeq z$ and

$Z(z) \simeq \beta(z) \simeq 1$. This approximation is equivalent to keeping in Eqs.(19) the terms of the order e^2 only:

$$\begin{aligned}\phi'(z) &\simeq -\frac{e^2\mu^\varepsilon}{2\pi^2\varepsilon}, \\ z\beta'(z) &\simeq \frac{e^2\mu^\varepsilon}{6\pi^2}, \\ zZ'(z) &\simeq \frac{e^2\mu^\varepsilon}{8\pi^2}\end{aligned}\tag{20}$$

in the Feynman gauge ($\alpha = 1$). We see that if we make the identification $z/z_0 = \mu_0/\mu$, where μ_0 is some IR scale smaller than μ , we find the well-known one-loop results [8], since we have then $z\partial_z = -\mu\partial_\mu$ (a decrease of z starting from z_0 is equivalent to an increase of μ starting from μ_0). One can understand this by noting that the decrease $zm_0 \rightarrow (z + \delta z)m_0$ of the electron mass influences mainly the propagation with momentum $p \approx zm_0$. In such a manner the modes whose dynamics is included in the effective action as z decreases ($dz < 0$) are approximately those which are included in the traditional RG step when the momentum scale of the cut-off is increased from $p = (z + \delta z)m_0$ to $p = zm_0$. This picture is lost for 1PI graphs containing two or more loops since the momenta of the internal loops appear as another scale which modifies the effect of the changing regulator parameters in a more complicated way. In this case our scheme will be different from the usual one, reflecting the different organization of the gradient and loop expansions.

Landau pole: It is interesting to note that the integration of equation (20),

$$\beta(z) \simeq 1 + \frac{e^2}{6\pi^2} \ln(z/z_0)\tag{21}$$

reproduces the well-known one-loop value of the Landau-pole,

$$\frac{z}{z_0} = \frac{\mu_0}{\mu} = \exp\left(-\frac{6\pi^2}{e^2}\right).\tag{22}$$

But the essential difference is that while the traditional Landau-pole is obstructing the further increase of the UV cut-off, the pole in the present scheme prevents us to further decrease the bare mass and indicates some problem with large amplitude quantum fluctuations. Since the initial condition of our evolution equation is already an effective potential corresponding to the regulated theory with a fixed cut-off, the UV Landau pole cannot be seen in our

scheme. In both cases the pole reflects the problem that the loop corrections dominate the tree-level part of the effective action when 'too many' modes are treated in the one-loop approximations.

But this similarity is an artifact of the autonomous solution of the truncated equation (20). Instead one should allow the other parameters, at least the fermion mass, to run as well. For this end we write the evolution equation of β as

$$\frac{1}{\beta'(z)} = \frac{6\pi^2}{e^2\mu^\varepsilon}(z + \phi(z)), \quad (23)$$

which gives

$$\frac{\beta''(z)}{\beta'(z)} = \left(\frac{3}{4\alpha\varepsilon} - \frac{6\pi^2}{e^2\mu^\varepsilon} \right) \beta'(z) + \frac{9}{4\varepsilon} \frac{\beta'(z)}{\beta(z)}, \quad (24)$$

after using the first equation of (19). It is easy to integrate this equation and find by means of (19) again:

$$\beta(z) = \left(\frac{z_0 + \phi(z_0)}{z + \phi(z)} \right)^{4\varepsilon/9} \exp \left\{ \frac{1 - \beta(z)}{3} \left(\frac{1}{\alpha} - \frac{8\pi^2\varepsilon}{e^2\mu^\varepsilon} \right) \right\} \quad (25)$$

It should be noted that we cannot naïvely take the limit $\varepsilon \rightarrow 0$ in the expression (25) since $\phi(z)$ is then diverging. This equation shows that $\beta(z)$ is non-vanishing for any value of z and therefore this pole is actually avoided by the evolution of the fermion mass and there is no problem to reconstruct the physical theory at $z = 0$. It is remarkable that the mechanism by which this happens, a running fermion mass, is actually the same how the Landau pole is claimed to be avoided in lattice QED by spontaneous symmetry breakdown of the chiral symmetry [9].

Conclusion: To conclude, we stress the analogies between the momentum dependence traced by the usual renormalization procedures and the dependence on the amplitude of quantum fluctuations. We showed that the leading order scaling law corresponding to small amplitudes (i) agrees with the traditional scaling law of the U.V. regime and (ii) is independent of the choice of the gauge as shown by Eqs.(17). The present work is the first step towards an extension of the renormalization scheme based on the field amplitude for non-Abelian gauge theories where a suitable gradient expansion should provide us with a deeper insight into the dynamics of the I.R. regime.

Acknowledgement: This work has been supported in parts by the TMR E.U. contract FMRX-CT97-0122, the Leverhulme Trust, U.K., the grants OTKA T29927/98, OTKA T032501/00, and DFG-MTA 436UNG113/140/0.

References

- [1] S.Coleman, E.Weinberg, Phys.Rev.D7: 1888 (1973).
- [2] F.J.Wegner, A.Houghton, Phys.Rev A8: 401 (1973).
- [3] J.M.Carmona, J.Polonyi, A.Tarancon, Phys.Rev.D61: 085018 (2000);
J.Alexandre, V.Branchina, J.Polonyi, Phys.Lett.B445: 351 (1999);
S.B.Liao, J.Polonyi, Phys.Rev.D51: 4474(1995).
- [4] J.Polchinski, Nucl.Phys.B231: 269 (1984).
- [5] C.Wetterich, Phys.Lett.B301: 90 (1993); M.Reuter, C.Wetterich,
Nucl.Phys.B391, 147 (1993); T.Morris, Int.J.Mod.Phys.A9: 2411
(1994); U.Ellwanger, Phys.Lett.B335, 364 (1994); J.Bergers, N.Tetradis,
C.Wetterich, hep-ph/0005122; S.Arnone, Yu.A.Kubyshin T.R.Morris,
K.Yoshida, J.Mod.Phys.A16: 1809 (2001); U.Ellwanger, Z.Phys.C76:
721 (1997); M. Simionato, Int. J. Math. Phys. A15:2121(2000); 2153
(2000).
- [6] J. Alexandre, J. Polonyi, Ann.Phys.286: 1 (2000).
- [7] J. Alexandre, Phys.Rev.D64: 045011 (2001).
- [8] S.Pokorski, *Gauge Field Theories*, Cambridge Monographs on Mathe-
matical Physics (1987), section 5-2.
- [9] M.Göckler, R.Horsley, V.Linke, P.Rackow, G.Shierholtz, H.Stüben,
Phys.Rev.Lett.80: 4119 (1998).